



US005851988A

United States Patent [19]**Sportsman et al.**[11] **Patent Number:** **5,851,988**[45] **Date of Patent:** **Dec. 22, 1998****[54] NONPEPTIDE INSULIN RECEPTOR AGONISTS**

[75] **Inventors:** Richard Sportsman, San Francisco;
Hugo O. Villar, Newark; Lawrence M. Kauvar, San Francisco; Wayne R. Spevak, Albany, all of Calif.

[73] **Assignee:** Terrapin Technologies, Inc., South San Francisco, Calif.

[21] **Appl. No.:** 784,854

[22] **Filed:** Jan. 15, 1997

[51] **Int. Cl.⁶** A61K 31/05; A61K 31/10;
A61K 31/17; A61K 31/655

[52] **U.S. Cl.** 514/4; 514/150; 514/477;
514/483; 514/562; 514/563; 514/596; 514/597;
514/598; 514/617; 514/618; 514/619; 514/708;
514/762; 514/866

[58] **Field of Search** 435/7.2, 18; 514/763,
514/150, 477, 483, 562, 563, 596, 597,
598, 617, 618, 619, 708, 762, 866, 4; 562/427

[56] References Cited**FOREIGN PATENT DOCUMENTS**

0172427 2/1986 European Pat. Off.

OTHER PUBLICATIONS

Brinkworth, R.I. et al. "Non-Peptide Anti-AIDS Agents: Inhibition of HIV-1 proteinase by Disulfonates," *Biochemical and Biophysical Research Communications* (1992) 188(2):624-630.

Kirchberger, J. et al. "Studies of the interaction of NADH oxidase from *Thermus thermophilis* HB8 with triazine dyes," *Journal of Chromatography A* (1994) 668:153-164.

Kletzien, R.E. et al. "Enhancement of Adipocyte Differentiation by an Insulin-Sensitizing Agent," *Molecular Pharmacology* (1991) 41:393-398.


Kobayashi, M. et al. "Pioglitazone Increases Insulin Sensitivity by Activating Insulin Receptor Kinase," *Diabetes* (1992) 41:476-483.

Primary Examiner—David L. Fitzgerald

Assistant Examiner—Michael Pak

[57] ABSTRACT

Modulation of the activity of the insulin receptor, enhancement of glucose uptake by cells, and other effects significant in the control and management of diabetes are accomplished using compounds of the formula



wherein

each A is independently a proton-accepting substituent;

each R is independently a noninterfering substituent;

n is 0, 1, or 2; and

each linker is independently an isostere of —NHCONH— or of —N=N— or of —NHCO—.

Compounds in the genus of Formula (1) can also be used for structure activity studies to identify features responsible for the relevant activities.

25 Claims, 9 Drawing Sheets

